

CARDIABET CRITICARE

S.No	PRODUCT NAME	COMPOSITION	PACKING	OLD MRP/BOX	NEW MRP/BOX	PACK
CARDIAC RANGE						
1	ATOLVA-10	Atorvastatin 10 mg 1. Reduces the risk of myocardial infarction. 2. Atorvastatin is an oral drug that lowers the level of cholesterol in the blood. 3. Atorvastatin also reduces the concentration of triglycerides in the blood.	ALU-ALU	500.00	468.75	10x10
2	ATOLVA-20	Atorvastatin 20 mg 1. Reduces the risk of myocardial infarction. 2. Atorvastatin is an oral drug that lowers the level of cholesterol in the blood. 3. Atorvastatin also reduces the concentration of triglycerides in the blood.	ALU-ALU	1200.00	1125.00	10x10
3	ATOLVA-C10	Atorvastatin 10 mg + Clopidogrel 75 mg 1. Effectively lowers plasma cholesterol & lipoprotein levels. 2. To prevent heart attacks & strokes in persons with heart disease (recent heart attack), recent stroke, or blood circulation disease (peripheral vascular disease).	ALU-ALU	950.00	970.00	10x10
4	ATOLVA-F	Atorvastatin 10mg + Fenofibrate 160mg 1. Atorvastatin is used along with a proper diet to help lower "bad" cholesterol and fats (such as LDL, triglycerides) and raise "good" cholesterol (HDL) in the blood. 2. Fenofibrate helps reduce cholesterol and triglycerides (fatty acids) in the blood. 3. High levels of these types of fat in the blood are associated with an increased risk of atherosclerosis (clogged arteries).	10x10 TAB.	1800.00	1687.50	10*10
5	AMTOPROL-AL	Metoprolol 50mg XL + Amlodipine 5mg 1. well documented efficacy in angina pectoris reducing the frequency, duration & severity of angina attacks. 2. A selective calcium channel blocker vaso dilator. 3. Reduces total peripheral resistance.	BLISTER	750.00	703.13	10x10
6	CALROV-10	Rosuvastatin Calcium 10mg 1. 54% relative risk reduction in myocardial infarction. 2. 48% relative risk reduction in stroke. 3. 47% relative risk reduction in revascularization procedures.	ALU-ALU	1050.00	984.38	10x10
7	CALROV-20	Rosuvastatin Calcium 20mg	ALU-ALU	1380.00	1293.75	10x10
8	CALROV-F	Rosuvastatin 10mg + Fenofibrate 160mg	ALU-ALU	1490.00	1396.88	10x10
9	TELISKA-40	Telmisartan HCL 40mg 1. Protective benefits against the vascular and renal damage caused by diabetes and cardiovascular disease. 2. Blocks angiotensin receptor.	ALU-ALU	950.00	950.00	10x15

10	TELISKA-80	Telmisartan HCL 80mg 1. Protective benefits against the vascular and renal damage caused by diabetes and cardiovascular disease. 2. Blocks angiotensin receptor.	ALU-ALU	1180.00	1106.25	10x15
11	TELISKA-H	Telmisartan HCL 40mg + Hydrochlorothiazide 12.5mg 1. Reduces arterial blood pressure. 2. Low dose thiazide diuretics are associated with an additional reduction in CHD & total mortality.	ALU-ALU	700.00	656.25	10x10
12	TELSIKA-AM	Telmisartan 40mg + Amlodipine 5mg 1. More effective than Losartan in controlling SBP. 2. Offers highest PPAR Gamma activation as compared to any other ARB	ALU-ALU	750.00	700.00	10x10
13	TELISKA-AM-H	Amlodipine 5mg + Telmisartan HCL 40mg + Hydrochlorothiazide 12.5mg 1. Reduces arterial blood pressure. 2. Low dose thiazide diuretics are associated with an additional reduction in CHD & total mortality. 3. More effective than Losartan in controlling SBP. 4. Offers highest PPAR Gamma activation as compared to any other ARB.	BLISTER	1300.00	1218.75	10x10
14	TELISKA-CL	Telmisartan 40mg+Clindipine 10mg	ALU-ALU	680.00	750.00	10x10
15	TELISKA-CM	Telmisartan 40mg+Clindipine 10mg+Metoprolol Succinate 50 mg	ALU-ALU		2450.00	10x10
16	TELISKA-MT25	Telmisartan 40mg+Metoprolol 25mg	10x10x ALU ALU	1450.00	1359.38	10x10
17	TELISKA-MT50	Telmisartan 40mg+Metoprolol 50mg	10x10x ALU ALU	1650.00	1546.88	10x10

DAIBETIC RANGE

18	EMPAFORIN 10	Empagliflozin 10mg Tablet	10x10 Blister	0.00	1350.00	10x10
19	EMPAFORIN 25	Empagliflozin 25g Tablet	10x10 Blister	0.00	2250.00	10x10
20	EMPAFORIN LN 25/5	Empagliflozin 25mg+Linagliptin 5mg	10x10 Blister	0.00	2450.00	10x10
21	SITAZLIN-M	DAPAGLIFLOZIN 10MG+SITAGLIPTON 100GM+METFORMIN SR 500MG	10x10 ALU ALU Bilayered		2250.00	10x10
22	GLISPO-1M	Glimepiride 1mg + Metformin 500mg (SR) 1. Highly effective in patients having both high fasting and high post-prandial blood glucose levels. 2. The addition of glimepiride to MET is an effective treatment for patients inadequately controlled on sulfonylurea or Met alone.	BLISTER (BI-LAYERED)	480.00	450.00	10x10
23	GLISPO-2M	Glimepiride 2mg + Metformin 500mg (SR) After the intensive treatment, FPG, 2 HPG & HBA1c decreased significantly (p>0.01); HOMA-IR decreased & HOMA-B increased significantly (p<0.01) & TG, CHOL, LDL decreased significantly (p<0.01) after treatment	BLISTER (BI-LAYERED)	720.00	675.00	10x10

24	GLISPO-2M FORTE	Glimepride 2mg + Metformin 1000mg SR (SR) After the intensive treatment, FPG, 2 HPG & HBA1c decreased significantly ($p > 0.01$); HOMA-IR decreased & HOMA-B increased significantly ($p < 0.01$) & TG, CHOL, LDL decreased significantly ($p < 0.01$) after treatment.	BLISTER (BI-LAYERED)	985.00	923.44	10x10
25	GLISPO-PM1/15	Glimepride 1mg + Metformin 500mg (SR) + Pioglitazone 15 mg 1. Stimulates the release of insulin from functioning pancreatic beta cells. 2. Counters insulin resistance. 3. decreases hepatic glucose production also decreases intestinal absorption of glucose.	BLISTER (BI-LAYERED)	640.00	600.00	10x10
26	GLISPO-PM2/15	Glimepride 2mg + Metformin 500mg (SR) + Pioglitazone 15 mg 1. Stimulates the release of insulin from functioning pancreatic beta cells. 2. Counters insulin resistance. 3. decreases hepatic glucose production also decreases intestinal absorption of glucose.	BLISTER (BI-LAYERED)	910.00	853.13	10x10
27	GLISPO-PM2/15	Glimepride 2mg + Metformin 500mg (SR) + Pioglitazone 15 mg 1. Stimulates the release of insulin from functioning pancreatic beta cells. 2. Counters insulin resistance. 3. decreases hepatic glucose production also decreases intestinal absorption of glucose.	BLISTER (BI-LAYERED)	1360.00	1270.00	10x15
28	TRIBOG-1.2	Glimepride 1mg + Voglibose 0.2mg + Metformin 500mg (SR) 1. Voglibose scores over both acarbose and miglitol in terms of side effect profile. 2. proven efficacy. 3. metformin causes few adverse effects the most common is gastrointestinal upset and is associated with a low risk of hypoglycemia.	BLISTER (BI-LAYERED)	940.00	881.25	10x10
29	TRIBOG-2.2	Glimepride 2mg + Voglibose 0.2mg + Metformin 500mg (SR) 1. Voglibose scores over both acarbose and miglitol in terms of side effect profile. 2. proven efficacy. 3. metformin causes few adverse effects the most common is gastrointestinal upset and is associated	BLISTER (BI-LAYERED)	1200.00	1125.00	10x10
30	TRIBOG-2.2	Glimepride 2mg + Voglibose 0.2mg + Metformin 500mg (SR) 1. Voglibose scores over both acarbose and miglitol in terms of side effect profile. 2. proven efficacy. 3. metformin causes few adverse effects the most common is gastrointestinal upset and is associated	BLISTER (BI-LAYERED)	900.00	1200.00	10x10
31	GLYCLIP-20	Teneligliptin 20 mg Teneligliptin is used in the treatment of type 2 diabetes It is used in addition to diet and exercise to improve blood sugar control in adults with type 2 diabetes	ALU-ALU	1250.00	1171.88	10x10
32	GLYCLIP-M	Telegliptin 20mg+Metformin 500mg	ALU-ALU	0.00	1620.00	10x10
33	LORIPTA-V	Vidagliptin 50mg It helps to control the levels of sugar (glucose) in your blood. If your body does not make enough insulin, or if it does not use the insulin it makes effectively, this results in the condition called sugar diabetes	ALU-ALU	950.00	890.63	10*10

34	LORIPTA-VM	Vidagliptin 50mg + Metformin 500 mg It is used to treat type 2 diabetes mellitus in people who are already taking vildagliptin and metformin tablets separately, or whose diabetes cannot be controlled by metformin alone Type 2 diabetes mellitus used to be known as 'non-insulin-dependent diabetes mellitus (NIDDM)' or 'maturity onset diabetes	BLISTER (BI-LAYERED)	1400.00	1312.50	10*10
35	VOBTRI-0.3MD	Voglibose 0.3mg (MD) Voglibose inhibitors delay glucose absorption at the intestine level and thereby prevent sudden surge of flucoose after a meal.	ALU-ALU	880.00	825.00	10x10